



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Metabolic model for Tetrasphaera involved in enhanced biological phosphorus removal plants

Kristiansen, Rikke; Nguyen, Hien Thi Thu; Saunders, Aaron Marc; Nielsen, Jeppe Lund; Wimmer, Reinhard; Le, Vang; Petrovski, Steve; McIlroy, Simon Jon; Seviour, Robert; Calteau, Alexandra; Nielsen, Kåre Lehmann; Nielsen, Per Halkjær

Publication date:
2012

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Kristiansen, R., Nguyen, H. T. T., Saunders, A. M., Nielsen, J. L., Wimmer, R., Le, V., Petrovski, S., McIlroy, S. J., Seviour, R., Calteau, A., Nielsen, K. L., & Nielsen, P. H. (2012). *Metabolic model for Tetrasphaera involved in enhanced biological phosphorus removal plants*. Poster presented at 7th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Metabolic Model for *Tetrasphaera* in Enhanced Biological Phosphorus Removal Plants

Rikke Kristiansen,¹ Hien T. T. Nguyen,¹ Aaron M. Saunders,¹ Jeppe L. Nielsen,¹ Reinhard Wimmer,¹ Vang Quy Le,¹ Steve Petrovski,² Simon J. McIlroy,^{1,2} Robert Seviour,² Alexandra Calteau,³ Kåre L. Nielsen,¹ and Per H. Nielsen¹

¹Department of Biotechnology, Chemistry and Environmental Engineering, Sohngaardsholmsvej 49, 9000 Aalborg, Denmark

²Biotechnology Research Centre, La Trobe University, Bendigo, Vic 3552, Australia

³LABGeM, CEA/DSV/IG/Genoscope, EVRY Cedex, France



INTRODUCTION

Enhanced biological phosphorus removal (EBPR) is an environmentally friendly and cost-effective phosphorus removal process for wastewater. It is believed to be carried out mainly by certain polyphosphate accumulating organisms (PAOs) called *Candidatus Accumulibacter*. Recently, actinobacteria belonging to the genus *Tetrasphaera* have been identified as putative polyphosphate accumulating organisms. New investigations show that they comprise a larger fraction of the biomass in full-scale Danish wastewater treatment plants than *Candidatus Accumulibacter* (8-12% and 3-7%, respectively) and may therefore be more important in the phosphorus removal process. Little is known about the diversity and physiology of *Tetrasphaera* but preliminary results indicate that their ecological niche is different from *Candidatus Accumulibacter*.

AIM

- Determine the diversity and abundance of *Tetrasphaera* in Danish treatment plants with enhanced biological phosphorus removal (EBPR).
- Develop a metabolic model for phosphate-removing *Tetrasphaera* based on genomic investigations and pure culture studies.

METHODS

Identification and quantification

Quantitative FISH using published and novel species-specific probes

Development of metabolic model

- Paired-end genome sequencing of four *Tetrasphaera* isolates using the Illumina platform.
- Annotation using MaGe (Microbial Genome Annotation Platform) from MicroScope.
- Validating the genomic findings with pure culture studies.

CONCLUSIONS

- A high abundance and diversity of *Tetrasphaera* were identified in Danish full-scale EBPR plants.
- Tetrasphaera* are potentially novel and ecologically relevant polyphosphate accumulating organisms in full-scale EBPR plants with a different physiology in EBPR plants compared to *Candidatus Accumulibacter*.

Identification and quantification

Fig. 1: Phylogenetic tree of 16S rRNA gene sequences of *Tetrasphaera* from EBPR plants. The sequences group in 3 clusters. 2 clusters include isolated *Tetrasphaera*.





Fig. 2: FISH images of *Tetrasphaera*.



Probe	Cluster	Treatment Plant (% EUBmix ± SD)				
		Ejby Mølle	Hjørring	Åbenrå	Skive	Odense
Elo1250	1	0	0	< 1	3.2 ± 1.3	1.2 ± 1.0
Tet823	1	< 1	3.4 ± 1.6	0	0	0
Tet266	1	5.5 ± 2.0	4.2 ± 1.2	4.1 ± 1.4	< 1	4.3 ± 1.0
Tet842	2	< 1	1.1 ± 1.0	3 ± 1.3	< 1	< 1
Tet831	2	0	0	3.1 ± 1.5	0	0
Tet892	2	5.4 ± 1.3	4.4 ± 1.0	4.5 ± 1.5	4.3 ± 1.0	4.4 ± 1.0
Tet87	2	2.3 ± 1.7	1.2 ± 1	< 1	< 1	< 1
Tet174	2	5.2 ± 1.3	4 ± 1.0	2.9 ± 1.0	1.7 ± 1.0	4.5 ± 1.2
Tet654	3	6.8 ± 2.0	5.7 ± 1.1	5.5 ± 1.8	5.4 ± 2.0	5.2 ± 1.4
Tet19	3	4.3 ± 1.5	4.1 ± 1.0	3.2 ± 1.0	3.5 ± 1.0	4.3 ± 1.0
PAO mix (<i>Accumulibacter</i>)		5.4 ± 1.0	3.1 ± 1.3	4.8 ± 1.5	4.5 ± 1.0	5.4 ± 1.2

Table 1: Abundance of *Tetrasphaera* in Danish EBPR plants determined using qFISH.

Development of metabolic model




Fig. 3: Simplified metabolic model of *Candidatus Accumulibacter* during the EBPR process (Martín et al., 2006).

Pathway	<i>Accumulibacter</i>	<i>T. australiensis</i>	<i>T. elongata</i>	<i>T. jenkinsii</i>	<i>T. japonica</i>
Glycolysis	+	+	+	+	+
Gluconeogenesis	+	+	+	+	+
Glycogen synthesis	+	+	+	+	+
Glycogenolysis	+	+	+	+	+
TCA cycle	+	+	+	+	+
Glyoxylate shunt	+	-	-	-	-
PHA synthesis	+	-	-	-	+
Polyphosphate metabolism	+	+	+	+	+
Fermentation	-	+	+	+	+
Acetate uptake	+	+	+	+	+
Glucose uptake	-	+	+	+	+
Glutamate/aspartate uptake	-	+	+	+	+

Table 2: Potential metabolic pathways identified by genomic investigations.

+: present
-: not present




Fig. 4: The ability of *T. elongata* to release Pi (triangles) and produce glycogen (diamonds) in the anaerobic phase feed with glucose (squares) (0-3 h) and take up Pi and consume glycogen in the aerobic phase (3.5-6.5 h).




Fig. 5: Metabolic model for *Tetrasphaera*. The key metabolic features enabling *Tetrasphaera* to compete in full-scale EBPR plants are shown.